

ACTIMAX

FERTILISER

Precision
technologies



N	P	K	Mg	S	Ca
58			9	81.6	67.14
B	Cu	Fe	Mn	Mo	Zn
44			80	3.9	



In g/L for liquid products

Specifications

PACK SIZE

10 L

FORMULATION

Specific Mineral Balance

CHECK CONDITION OF PRODUCT AT

24 months

APPLICATION

Foliar

FORM

Liquid (SC)

DENSITY

1.455

STORAGE CONDITIONS

Store away from frost.

Agronomic interests

ROLE OF NUTRIENT(S)

Magnesium : Photosynthesis / Synthesis of carbohydrates and lipids

Boron : Role on flowering and fruit set / Migration on sugars / Protein synthesis / Meristematic activity and growth / Use of auxin in the plant

Molybdenum : Nitrate reduction / Protein synthesis / Contributes to the nodulation of legumes

Manganese : Enzyme activation / Chlorophyll synthesis / Nitrate reduction and protein synthesis

Sulphur : Protein constituents / Enzymatic role / Photosynthesis

Calcium: Improvement in yield quality / role in meristematic activity

PRODUCT SPECIFICATIONS

Actimax prevents and corrects the deficiencies in Boron, Calcium, Molybdenum, Sulphur and Manganese of all crops by foliar application.

The rigorous selection of the raw materials guarantees it to be 100 % soluble in the treatment mix.

FEATURES AND BENEFITS OF THE FORMULATION

The unique formulation of this product guarantees maximum foliar uptake, without the risk of phytotoxicity, designed to achieve nutrient absorption in less than 6 hours.

This product is formulated with rigorously selected raw materials, guaranteed by quality control, to ensure a perfect safety of use and to optimize the absorption through the leaves.

Directions of use

CROP RECOMMENDATIONS

Crops	Rate L/ha	Water Volume L /ha	Timing
Beet	3-5 L	150 L min	2 applications during starting period and on well established crops
Field brassicas Oil Seed, Rape			1-2 applications from 2-3 leaves to start of flowering
Broccoli, Cabbage, Cauliflower ...			From stage 4-6 leaves (2 weeks after transplanting) and repeat application after 15 days
Peas and Legumes			1st Application at Pre - Flowering followed by a 2nd application at Flat Pod stage.
Lettuce			From stage 6-8 leaves

Instructions for use

For best results, use sufficient water to achieve an even crop coverage to the point of run-off.

Avoid spraying in very strong sunlight and/or very high temperatures. If possible, apply during evening or early morning especially if day-time temperatures exceed 28°C. Very young foliage and crops under stress may be more susceptible to scorch. Lower doses should be applied in these circumstances and the application interval reduced accordingly. Refer to the Crop Recommendations table for application rates and timings for individual crops.

Precautions

MIXING

1. Fill the spray tank with half the required amount of water and start agitation.
2. Add the required amount of Actimax and then add the rest of the water, maintaining agitation continuously. When combining with other components in a tank-mix, consult the manufacturer and distributor concerned. Do not allow the mixture to stand without agitation.
3. After spraying, clean and rinse the spraying equipment thoroughly.

IMPORTANT: respect the uses, doses, conditions and instructions for use mentioned on the packaging, which are determined according to the characteristics of the product and the applications for which it is recommended. On this basis, conduct the crop and treatments according to the best agricultural practices taking into account, under your responsibility, all the specific factors concerning your farm, such as soil characteristics, weather conditions, cropping methods, cultivars and their specific resistances. AGRONUTRITION guarantees the quality its products only in their initial packaging over the period indicated on the manufacturer's certificates. They guarantee their conformity to the composition indicated on the packaging and to the regulations currently in force. Precautionary measure brochures are available on request. **To request a MSDS, thank you to contact: fds-msds@agro-nutrition.fr**