

FERTIGONIA 13-18-11

NPK FERTILISER 13-18-11 WITH MICRO-NUTRIENTS
FERTILISER OBTAINED BY BLENDING

Agro nutrition

Precision
technologies

N	P	K	Mg	S	Ca
13%	18%	11%	0.3%		
B	Cu	Fe	Mn	Mo	Zn
0.01%	0.008%	0.026%	0.032%		0.023%



Cu, Fe, Mn and Zn are chelated by EDTA

Specifications

PACK SIZE

25 KG

FORMULATION

Specific Mineral balance

CHECK CONDITION OF PRODUCT AT

36 months

APPLICATION

Foliar - Soil

FORM

Powder (SP)

DENSITY

1

STORAGE CONDITIONS

Store away from humidity.

Agronomic interests

ROLE OF NUTRIENT(S)

Supply combined with Nitrogen, Phosphorus, Potassium, Magnesium and Trace elements essential to the good progress of the crop. The proportion of each element is specific to the needs of the plant then allowing an improvement of the overall nutritional status of the plan

PRODUCT SPECIFICATIONS

FERTIGONIA is a polyvalent product which is suitable to numerous crops. It is specifically adapted to nutrition, improvement and growth of the vegetative system

FEATURES AND BENEFITS OF THE FORMULATION

This product is formulated with rigorously selected raw materials, guaranteed by quality control, to ensure a perfect safety of use and to optimize optimise both foliar and root absorption.

Directions of use

CROP RECOMMENDATIONS

May be applied to all crops in the generative growth/fruiting stage when necessary to ensure the plant is at a balanced nutrient equilibrium level.

SOIL Applications - Fertigation - all crops: 1-3 g/L at 10-14 day intervals.

	1 g/L	2 g/L	3 g/L
ECmS/cm	1.181	2.019	2.942
pH	5.20	5.12	5.08

Acrecio and Fertigonía 13-18-11 can be applied together through fertigation.

FOLIAR Applications

/ Cereals and Row Crops: 2Kg - 3Kg/Ha with a minimum of 150 - 300L water rate at 10-14 day intervals.

/ Citrus, Avocado, 3-5kg/ha

min 400-1000L water/ha at 10-14 day intervals

/ Grapes 3-5kg/ha min 400-1000L water/ha at 10-14 day intervals.

/ Other crops: 2-5kg/ha in a minimum 400-1000L water/ha at 10-14 day intervals.

Instructions for use

For best results, use sufficient water to achieve an even crop coverage to the point of run-off. Avoid spraying in very strong sunlight and/or very high temperatures. If possible, apply during evening or early morning especially if day-time temperatures exceed 28°C. Very young foliage and crops under stress may be more susceptible to scorch. Lower doses should be applied in these circumstances and the application interval reduced accordingly. Refer to the Crop Recommendations table for application rates and timings for individual crops.

Precautions

COMPATIBILITY: Always read the product labels and follow the manufacturers' instructions for all products at all times. Some settings, outside the control of the manufacturer or distributor, may have repercussions on the performance of co-applied products. Therefore, the co-application is made at the risk of the end user. When combining with other components in a tank-mix, always add this product first. If there is any doubt at all, consult the manufacturer or distributor concerned.

IMPORTANT: respect the uses, doses, conditions and instructions for use mentioned on the packaging, which are determined according to the characteristics of the product and the applications for which it is recommended. On this basis, conduct the crop and treatments according to the best agricultural practices taking into account, under your responsibility, all the specific factors concerning your farm, such as soil characteristics, weather conditions, cropping methods, cultivars and their specific resistances. AGRONUTRITION guarantees the quality its products only in their initial packaging over the period indicated on the manufacturer's certificates. They guarantee their conformity to the composition indicated on the packaging and to the regulations currently in force. Precautionary measure brochures are available on request. **To request a MSDS, thank you to contact: fds-msds@agro-nutrition.fr**

